

Secondary Math 2H

Unit 10 – Probability

10.1 The Fundamental Counting Principle / Factorial notation / Probability

Introduction to Factorial Notation Factorial - the product of an integer and all integers below it.

Used for - how many ways you can arrange / or order a set

Example: There are 8 people running a race. How many different outcomes for the race are there?

Solution: There are 8 different people who can finish first. Once someone finishes first, there are only 7 people left competing for second place, then six left competing for third, and so on.

So, to calculate all the different outcomes for the race:

$$8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 40,320 \quad \text{or} \quad 8!$$

There are 40,320 different outcomes for the race.

8 7 6 5 4 3 2 1

The example above requires you to multiply a series of descending natural numbers: $8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$. This can be written as $8!$ (read 8 factorial).

$8!$ means $8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$

$$5! = 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 120$$

$$9! = 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = 362,880$$

It is generally accepted that $0! = 1$.

Now, what if you had to calculate $20!$? Do you want to enter all of those numbers into your calculator?

The factorial key on your calculator can be found by:

math
Prob
 $\frac{4}{\square} \Rightarrow !$

Examples: Simplify each expression to a single number or fraction.

Show work

$$1. \frac{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} = \boxed{6}$$

$$2. \frac{8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1} = \boxed{56}$$

$$3. 3! = 3 \cdot 2 \cdot 1 = \boxed{6}$$

$$4. 6! = 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 = \boxed{720}$$

$$5. \frac{4!}{3!} = \frac{4 \cdot 3 \cdot 2 \cdot 1}{3 \cdot 2 \cdot 1} = \boxed{4}$$

$$6. \frac{6!}{4!} = \frac{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{1 \cdot 2 \cdot 3 \cdot 2 \cdot 1}$$

$$= \boxed{30}$$

$$7. \frac{101!}{99!} = \frac{101 \cdot 100 \cdot 99!}{99!} = \boxed{10,100}$$

OR

$$\frac{65 \cdot 4!}{4!} = \boxed{30}$$

Fundamental Counting Principle:

If event M can occur in m ways and event N can occur in n ways, then event M followed by event N can occur in $m \cdot n$ ways.

If you are making a sandwich and there are three types of bread, five types of meat, and four types of cheese available and you choose one type of bread, one type of meat, and one type of cheese, there are $3 \cdot 5 \cdot 4 = 60$ different sandwiches you can make.

Examples:

- a) How many possible locker combinations are available if each combination consists of three numbers between 1 and 50?

$$\underline{50} \cdot \underline{50} \cdot \underline{50} = \boxed{125,000}$$

- b) How many license plates consisting of three numbers followed by three letters are possible if numbers and letters can be repeated?

10 single digit #s
0, 1, 2, 3, 4, 5, 6, 7, 8, 9

$$\underline{10} \cdot \underline{10} \cdot \underline{10} \cdot \underline{26} \cdot \underline{26} \cdot \underline{26} = \boxed{17,576,000}$$

- c) There are eight true-false questions on a quiz. How many different answer combinations are possible?

$$\underline{2} \cdot \underline{2} = \boxed{256}$$

- d) A golf club manufacturer makes drivers with 4 different shaft lengths, 3 different lofts, 2 different grips, and 2 different club head materials. How many different combinations are possible?

$$4 \cdot 3 \cdot 2 \cdot 2 = \boxed{48}$$

- e) For a college application, George must select one of five topics on which to write a short essay. He must select a different topic from the list for a longer essay. How many ways can he choose the topics for the two essays?

$$\underline{5} \cdot \underline{4} = \boxed{20}$$

- f) Abby is registering at a website. She must create a password containing 6 different numbers. How many possible passwords are there?

not same as
10

$$\underline{10} \underline{9} \underline{8} \underline{7} \underline{6} \underline{5} = \boxed{151,200}$$

- g) How many ways can 8 books be arranged on a shelf?

$$\underline{8} \underline{7} \underline{6} \underline{5} \underline{4} \underline{3} \underline{2} \underline{1} = 8! \\ \boxed{40,320}$$


Probability:

A value that represents the likelihood of an event. It can be expressed as a fraction, decimal, or a percentage. A probability of 0 means that the event is impossible and a probability of 1 (or 100%) means that the event is certain to occur.

$$\text{probability} = \frac{\text{total # of favorable outcomes in the category of interest}}{\text{total # of possible outcomes}}$$

Example:

A spinner is pictured below. If the arrow is spun, what is the probability that the spinner lands on:



- a) green $\frac{3}{12} = \boxed{\frac{1}{4}}$ or .25 or 25%
- b) yellow $\boxed{\frac{1}{12}}$ or .0833 or 8.3%
- c) blue or red $\boxed{\frac{8}{12}}$ or .666 or 66.7%
or .67
- d) not red $\boxed{\frac{7}{12}}$ or .583 or 58.3%

Example:

A card is drawn at random from a standard deck of cards. What is the probability that the card drawn is:

Sample Space for Choosing a Card from a Deck

Ace	2	3	4	5	6	7	8	9	10	Jack	Queen	King	13
♦	♦	♦	♦	♦	♦	♦	♦	♦	♦	♦	♦	♦	13
♠	♠	♠	♠	♠	♠	♠	♠	♠	♠	♠	♠	♠	13
♣	♣	♣	♣	♣	♣	♣	♣	♣	♣	♣	♣	♣	13

a) a heart $\frac{13}{52} = \boxed{\frac{1}{4}}$

b) red $\frac{26}{52} = \boxed{\frac{1}{2}}$

c) a 5 $\frac{4}{52} = \boxed{\frac{1}{13}}$

d) red and a 4 $\frac{2}{52} = \boxed{\frac{1}{26}}$

e) not an ace $\frac{48}{52} = \boxed{\frac{12}{13}}$

f) black or a king $\frac{26+2}{52} = \boxed{\frac{7}{13}}$

$$\frac{26}{52} = \boxed{\frac{7}{13}}$$

10.2 Permutations and Combinations

Independent and Dependent Events

VOCABULARY -

- **Outcome:** The result of a single trial, experiment, or decision.
 - The possible outcomes of a coin flip would be heads or tails.
 - The possible outcomes of rolling a single die would be 1, 2, 3, 4, 5, or 6.
- **Sample Space:** The set of all possible outcomes.
 - The sample space for a coin flip is {H, T}.
 - The sample space for two coin flips would be {HH, HT, TH, TT}
- **Event:** One or more possible outcomes of a trial.
 - A coin flip coming up heads
 - Rolling either a 2 or a 6 on a die
 - Choosing a face card from a deck of cards
- **Independent Events:** The outcome of one event does not affect the outcome of another event.
 - Two coin flips are independent because the outcome of the first coin flip does not affect the outcome of the second flip.
- **Dependent Events:** The outcome of one event does affect the outcome of another event.
 - Choosing a piece of candy from a jar and then choosing a second piece without replacing the first are dependent events because taking the first piece affects what is available to be taken next.

Examples: Decide whether the events are *dependent* or *independent*.

1. Choosing an ice cream flavor and then choosing a topping for the ice cream.

Independent

2. Choosing one book on which to write an essay on and then a different book on which to give a presentation.

Dependent

3. Awarding 1st, 2nd, and 3rd prize to entries in an art contest.

Dependent

4. Selecting a fiction book and a nonfiction book from the library.

Independent

Permutation and Combinations

- **Permutation:** An arrangement of a group of objects, where order matters.
Ex: Selecting a president, vice president, and secretary/lock combination/batting order

Permutation Formula (no repetition allowed)

$$\frac{n!}{r!(n-r)!} = \frac{n!}{(n-r)!r!}$$

where n is the number of things you choose from and you choose r of them

- **Combination:** An arrangement or selection of objects in which **order** is not important.
Ex: Choosing Pizza toppings/selecting people to a committee

Combination Formula (no repetition allowed)

$$\underline{\underline{n}} C_r = \underline{\underline{C(n,r)}} = \underline{\underline{\binom{n}{r}}} = \underline{\underline{\frac{n!}{r!(n-r)!}}} \quad \text{where } n \text{ is the number of things you choose from and you choose } r \text{ of them}$$

Examples: Determine whether each situation involves a permutation or a combination.

Permutation \Rightarrow order
 $n^P_r \text{ or } P(n, r) = \frac{n!}{(n-r)!}$

Combination \Rightarrow no order
 $nC_r \text{ or } C(n, r) \text{ or } \binom{n}{r} = \frac{n!}{r!(n-r)!}$

Examples: Evaluate each expression

1. ${}_6P_2 = \frac{6!}{(6-2)!} = \frac{6!}{4!} = \frac{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{4 \cdot 3 \cdot 2 \cdot 1}$
Show this work or $\frac{6 \cdot 5 \cdot 4!}{4!} = \boxed{30}$

2. $P(10, 5) = \frac{10!}{(10-5)!} = \frac{10!}{5!} = \frac{10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5!}{5!}$
 $\boxed{30,240}$

3. ${}_9P_6 = \frac{9!}{(9-6)!} = \frac{9!}{3!} = \frac{9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3!}{3!}$
 $= \boxed{60,480}$

4. $P(25, 20) = \frac{25!}{(25-20)!} = \frac{25 \cdot 24 \cdot 23 \cdot 22 \cdot 21 \cdot 20 \cdot 19!}{5!}$
calc. $\frac{25!}{5!} = 1.292600837 \times 10^{23}$ or $\boxed{1.292600837 \times 10^{23}}$

5. ${}_{15}C_4 = \frac{15!}{4!(15-4)!} = \frac{15 \cdot 14 \cdot 13 \cdot 12 \cdot 11!}{4 \cdot 3 \cdot 2 \cdot 1 \cdot 11!}$
 $= \frac{32760}{11!} = \boxed{1365}$

6. ${}_9C_6 = \frac{9!}{6!(9-6)!} = \frac{9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3!}{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 3!}$
 $= \frac{504}{6} = \boxed{84}$

7. $C(12, 4) = \frac{12!}{4!(12-4)!} = \frac{12 \cdot 11 \cdot 10 \cdot 9 \cdot 8!}{4 \cdot 3 \cdot 2 \cdot 1 \cdot 8!}$
 $= \boxed{495}$

Examples: Determine whether each situation involves a *permutation* or a *combination*. Then find the number of possibilities.

1. Twelve skiers are competing in the final round of the Olympic freestyle skiing aerial competition. In how many ways can 3 of the skiers finish first, second, and third to win the gold, silver, and bronze medals?

12 \cdot 11 \cdot 10 = Permutation $\boxed{1320}$

$12P_3 = \frac{12!}{(12-3)!} = \frac{12!}{9!} = \frac{12 \cdot 11 \cdot 10 \cdot 9!}{9!} = \boxed{1320}$

2. A pizza shop offers twelve different toppings. How many different three-topping pizzas can be formed with the twelve toppings?

no order
Combination

$12C_3 = \frac{12!}{3!(12-3)!} = \frac{12!}{3 \cdot 2 \cdot 1 \cdot 9!} = \boxed{220}$

3. Your English teacher has asked you to select 3 novels from a list of 10 to read as an independent project. In how many ways can you choose which books to read?

Combination

$10C_3 = \frac{10!}{3!(10-3)!} = \frac{10 \cdot 9 \cdot 8 \cdot 7!}{3 \cdot 2 \cdot 1 \cdot 7!} = \boxed{120}$

4. The school yearbook has an editor-in-chief and an assistant editor-in-chief. The staff of the yearbook has 15 students. How many different ways can students be chosen for these 2 positions?

order
15 \cdot 14 = 210

$15P_2 = \frac{15!}{(15-2)!} = \frac{15 \cdot 14 \cdot 13!}{13!} = \boxed{210}$

$$5. \text{ Combination lock} \Rightarrow \text{Permutation ordered}$$

$$\underline{10} \cdot \underline{9} \cdot \underline{8} = 720$$

It is important to note that when you use the permutation or combination formulas above, repetition is not allowed. In other words, you can't have the same person win first and second place.

Another Case to Consider

Example: How many different ways can the letters MATH be arranged to create four-letter "words"?

Solution: This is an example of a permutation because the order of the letters would produce a different "word" or outcome. So, we use the permutation formula:

$${}_4P_4 = P(4,4) = \frac{4!}{(4-4)!} = \frac{4!}{0!} = \frac{4 \cdot 3 \cdot 2 \cdot 1}{1} = 24$$

$$\underline{4} \underline{3} \underline{2} \underline{1} = 24$$

But, what if some of the letters repeated? For example, how many ways can the letters in CLASSES be rearranged to create 7 letter "words"? Since the letter S repeats 3 times, some of the permutations will be the same so we will have to eliminate them. Here is how we do it. There are 7 letters to choose from and we are choosing 7 of them, so we would have the following:

S repeats 3 times

$${}_7P_7 = P(7,7) = \frac{7!}{(7-7)!} = \frac{7!}{0!} = \frac{7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{1} = 5040$$

Hold on – this is not our answer yet. We have to divide out our duplicate letters. As we mentioned earlier, the letter S repeats 3 times so we divide our answer by 3!:

$$\frac{5040}{3!} = 840$$

* Divide by factorials for any repeating letters.

Example 7: How many ways can the letters in MISSISSIPPI be arranged to create 11-letter "words"?

I repeats 4 times

S repeats 4 times

P repeats 2 times

$$\frac{11!}{4! 4! 2!} = \boxed{34,650}$$

Example 8:
STREETS

$$\frac{7!}{2! 2! 2!} = \boxed{630}$$

Example 9:
ASSIGNMENTS

$$\frac{11!}{3! 2!} = \boxed{3,326,400}$$

10.3 Venn Diagrams

Sample Space: The set of all possible outcomes for a chance process.

Event/Subset: An outcome or set of outcomes from the sample space.


Complement (A^c): “Not”

- All outcomes in the sample space that are not part of the event.

Chance Process	Sample Space	Event/Subset	Complement
Flip a coin	$S = \{\text{heads, tails}\}$	$B = \{\text{heads}\}$	$B^c = \{\text{tails}\}$
Roll a die	$S = \{1, 2, 3, 4, 5, 6\}$	even numbers $E = \{2, 4, 6\}$	$E^c = \{1, 3, 5\}$
Pick a letter in the word “probability”	$S = \{P, R, O, B, A, I, L, T, Y\}$	vowels $V = \{O, A, I, Y\}$	$V^c = \{P, R, B, L, T\}$


Union ($A \cup B$): “Or”, “Either”

- All of the elements that are in A or B or both.



Intersection ($A \cap B$): “And”, “Both”, “Overlap”, “In common”


- All of the elements that are in *both* A and B .
- If the two sets don't have anything in common, the intersection is the “empty set”, indicated by \emptyset or $\{\}$.



Note: If you want to write “everything in A that isn’t in B ,” you can write either


$A \cap B^c$ or $A - B$

\uparrow subtract
“(take away)”




Examples: Shade the appropriate portion of the Venn diagram.


1. A^c



2. $(A \cap B)^c$



3. $B - A$



Everything not in A
 B^c

Everything not in A and B

Everything in B, but not in A

Examples:

- Chance Process: Rolling a 10-sided die.
 - Event A: Rolling an odd number
 - Event B: Rolling a prime number

- a. What is the sample space?

$$S = \{1, 2, 3, 4, 5, 6, 7, 8, 9, 10\}$$


1 is not prime
2 is only even prime

- b. List the outcomes in each event

ODD $A = \{1, 3, 5, 7, 9\}$

Prime $B = \{2, 3, 5, 7\}$

- c. Draw a Venn diagram representing the sample space with subsets A and B.



Always start with middle
(what's in both sets)

- d. List all the outcomes in $A \cup B$.

$$A \cup B = \{1, 2, 3, 5, 7, 9\}$$

and, 'intersection'

- e. List all the outcomes in $A \cap B$.

$$A \cap B = \{3, 7, 5\}$$

- f. List all the outcomes in A^c .

not in A $A^c = \{2, 4, 6, 8, 10\}$

- g. List all the outcomes in $(A \cup B)^c$.

not in A or B $(A \cup B)^c = \{4, 6, 8, 10\}$

- h. List all the outcomes in $A - B$.

A but not B $A - B = \{1, 9\}$


- Chance Process: Reaching into a messy refrigerator and grabbing a food at random.
- Sample Space: $S = \{\text{broccoli, carrots, moldy cheese, milk, orange, lettuce, lime jello, bologna, egg, corn, celery}\}$
 - Event A: Picking a vegetable
 - Event B: Picking something green

- a. List the outcomes in each event.

$$A = \{\text{broccoli, carrots, lettuce, corn, celery}\}$$

$$B = \{\text{moldy cheese, lime jello, broccoli, lettuce, celery}\}$$

- b. Draw a Venn diagram representing the sample space with subsets A and B.



- c. List all the outcomes in $A \cup B$.

$$A \cup B = \{\text{carrots, corn, broccoli, lettuce, celery, moldy cheese, lime jello}\}$$

- d. List all the outcomes in $A \cap B$.

$$A \cap B = \{\text{broccoli, lettuce, celery}\}$$

- e. List all the outcomes in B^c . not in set B

$$B^c = \{\text{carrots, corn, orange, egg, milk, bologna}\}$$

- f. List all the outcomes in $(A \cap B)^c$. not in A and B

$$(A \cap B)^c = \{\text{orange, egg, milk, bologna, moldy cheese, lime jello, carrots, corn}\}$$

- g. List all the outcomes in $B - A$. Everything in B but not in A

$$B - A = \{\text{moldy cheese, lime jello}\}$$

Examples:

A political ad was run on TV and on radio.

- 33% of people saw it on TV.
- 21% heard it on the radio.
- 10% of people both saw it on TV and heard it on the radio.

Determine what percent:

- a) only saw it

$$23\%$$

- b) only heard it

$$11\%$$

- c) neither heard it or saw it

$$56\%$$

- d) did not see it

$$11 + 56 = 67\%$$

e) didn't hear it

$$23 + 56 = 79\%$$

A sample of 60 people are asked if they enjoy watching basketball and if they enjoy watching football.

- 25 people say they enjoy watching football
- 40 people say they enjoy watching basketball
- 15 people say they enjoy watching both

Determine how many people:

- a) enjoy football but not basketball

$$10$$

- b) enjoy basketball but not football


$$25$$

- c) don't enjoy either basketball or football


$$10$$

- d) don't like football

$$25 + 10 = 35$$



60




$$60 - 10 - 15 - 25 = 10$$

10.4 Probabilities from Venn Diagrams


Probability: A value that represents the likelihood of an event. It can be expressed as a fraction, decimal, or a percentage. A probability of 0 means that the event is impossible and a probability of 1 (or 100%) means that the event is certain to occur.

$$\text{probability} = \frac{\text{total # of favorable outcomes in the category of interest}}{\text{total # of possible outcomes}}$$



Remember, $(A \cap B)$ means “*A and B*” and $(A \cup B)$ means “*A or B (or both)*”. With “or” probabilities, make sure you don’t count the individuals who fall in both categories twice!

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$



Example: In the Math Club, there are 34 students. Eleven of the students are seniors, including 7 of the 20 girls. A student is chosen at random from the club. Fill in the table and find the following probabilities: * write answer as reduced fraction .

a) $P(\text{boy}) = \frac{\text{boys}}{\text{TOT}} = \frac{14}{34} = \boxed{\frac{7}{17}}$

b) $P(\text{senior}) = \frac{\text{Sen}}{\text{TOT}} = \boxed{\frac{11}{34}}$

overlap and
c) $P(\text{boy} \cap \text{senior}) = \frac{4}{34} = \boxed{\frac{2}{17}}$

d) $P(\text{girl} \cup \text{non-senior})$ Formula

	Seniors	Non-Seniors	Total
Boys	4	10	14
Girls	7	13	20
Total	11	23	34

$$\frac{20+23-13}{34} = \frac{30}{34} = \frac{15}{17}$$

$$\frac{7+13+10}{34} = \frac{30}{34} = \frac{15}{17}$$

Example: The number of students in a high school is 1400. Of those students, 550 take French, 700 take algebra, and 400 take both French and algebra. Fill in the Venn diagram, then find the following probabilities.


* Write answers as percent rounded to nearest tenth.

a) $P(\text{does not take French}) = \frac{300+550}{1400} = \frac{850}{1400} = 60.7\%$

b) $P(\text{algebra} \cap \text{French}) = \frac{400}{1400} = 28.6\%$

c) $P(\text{algebra, but not French}) = \frac{300}{1400} = 21.4\%$

d) $P(\text{algebra} \cup \text{French}) = \frac{550+700-400}{1400} = \frac{850}{1400} = 60.7\%$



Conditional Probability: The probability of an event occurring when we already know that another event has occurred.

$\curvearrowleft \text{ given}$
Example: $P(\text{lung cancer} | \text{smoke})$ would mean the probability of a person getting lung cancer given that the person smokes.

Conditional Probability Formula: $P(A|B) = \frac{P(A \cap B)}{P(B)}$ or $\frac{\text{total # in } A \cap B}{\text{total # in } B}$

★ “And” and “or” probabilities are fractions of the entire sample, but with conditional probabilities, the condition becomes the denominator of the fraction!

Examples:

An ice cream shop keeps track of whether people order vanilla or chocolate ice cream and whether they order a sugar cone or a waffle cone. Fill in the table and find the requested probabilities.

	Sugar Cone	Waffle Cone	Total
Vanilla	35	26	61
Chocolate	51	47	98
Total	86	73	159

* Leave answers as decimals rounded to the nearest thousandth.

a) $P(\text{vanilla}) = \frac{61}{159} = .384$ b) $P(\text{waffle}) = \frac{73}{159} = .459$ c) $P(\text{sugar}) = \frac{86}{159} = .541$

d) $P(\text{chocolate}) = \frac{98}{159} = .616$ e) $P(\text{vanilla and sugar}) = \frac{35}{159} = .220$ f) $P(\text{vanilla or waffle}) = \frac{26}{159} = .164$

g) $P(\text{chocolate or sugar}) = \frac{51}{159} = .321$

h) $P(\text{chocolate or waffle}) = \frac{47}{159} = .296$

i) $P(\text{vanilla or sugar}) = \frac{35+51+26}{159} = \frac{112}{159} = .704$

j) $P(\text{vanilla or waffle}) = \frac{35+26+47}{159} = .679$

k) $P(\text{chocolate or sugar}) = \frac{35+51+47}{159} = .836$

l) $P(\text{chocolate or waffle}) = \frac{51+47+26}{159} = .780$

m) $P(\text{vanilla} | \text{sugar}) = \frac{35}{86} = .407$

n) $P(\text{vanilla} | \text{waffle}) = \frac{26}{73} = .356$

o) $P(\text{chocolate} | \text{sugar}) = \frac{51}{86} = .593$

p) $P(\text{chocolate} | \text{waffle}) = \frac{47}{73} = .677$

q) $P(\text{sugar} | \text{vanilla}) = \frac{35}{61} = .574$

r) $P(\text{sugar} | \text{chocolate}) = \frac{51}{98} = .520$

s) $P(\text{waffle} | \text{vanilla}) = \frac{26}{61} = .426$

t) $P(\text{waffle} | \text{chocolate}) = \frac{47}{98} = .480$


conditional probabilities

conditional Probabilities w/Venn Diagrams

Examples: Use the Venn diagram to find the following probabilities.
 *write answer as reduced fraction.

a) $P(\text{job}|\text{male}) = \frac{27}{48} = \boxed{\frac{9}{16}}$

b) $P(\text{female}|\text{job}) = \frac{33}{60} = \boxed{\frac{11}{20}}$



c) $P(\text{male}|\text{no job}) = \frac{21}{21+19} = \boxed{\frac{21}{40}}$

d) $P(\text{no job}|\text{female}) = \frac{19}{33+19} = \boxed{\frac{19}{52}}$

e) A student from the sample works at McTaco. What is the probability that the student is male?

$$P(\text{male}|\text{job}) = \frac{27}{60} = \boxed{\frac{9}{20}}$$

f) Is a student from the sample more likely to have a job if he is a male? Justify your answer using conditional probability.

$$P(\text{job}|\text{male}) = \frac{27}{48} = 56.3\%$$

$$P(\text{job}|\text{female}) = \frac{33}{33+19} = \frac{33}{52} = 63.5\%$$

Females more likely to have a job.

Examples: Real-estate ads suggest that 64% of homes have a garage, 21% have a pool, and 17% have both a garage and a pool. Fill in the Venn diagram, then answer the following questions.

*write answer as percent. Round to nearest percent. 100%.

a) Find $P(\text{garage} \cup \text{pool})$ or

$$\frac{47+17+4}{100} = \frac{68}{100} = 68\%$$

b) Find $P(\text{garage}|\text{pool})$

$$\frac{17}{17+4} = \frac{17}{21} = .809 \approx 81\%$$

c) Find $P(\text{pool}|\text{garage})$

$$\frac{17}{64} = 27\%$$

d) Find $P(\text{pool}|\text{no garage})$

$$\frac{4}{4+32} = \frac{4}{36} = 11\%$$

e) Find $P(\text{no pool}|\text{garage})$

$$\frac{47}{64} = 73\%$$

f) Find $P(\text{no garage}|\text{no pool})$

$$\frac{32}{47+32} = \frac{32}{79} = 41\%$$

